

Dimer statistics on the Möbius strip and the Klein bottle

W. T. Lu and F. Y. Wu
Department of Physics
Northeastern University
Boston, Massachusetts 02115, USA

February 1, 2008

Abstract

Closed-form expressions are obtained for the generating function of close-packed dimers on a $2M \times 2N$ simple quartic lattice embedded on a Möbius strip and a Klein bottle. Finite-size corrections are also analyzed and compared with those under cylindrical and free boundary conditions. Particularly, it is found that, for large lattices of the same size and with a square symmetry, the number of dimer configurations on a Möbius strip is 70.2% of that on a cylinder. We also establish two identities relating dimer generating functions for Möbius strips and cylinders.

1 Introduction

It is well-known that the problem of closed-packed dimers on planar graphs can be solved, meaning that its generating function can be formulated as a Pfaffian, the square root of an antisymmetric determinant [1]. This leads to the possibility that the generating function can be evaluated in a closed form for lattices with periodic structures. Indeed, Kasteleyn [1] has succeeded in obtaining the closed-form expressions of the generating function for the simple quartic lattice with both free and toroidal boundary conditions, and McCoy and T. T. Wu [2] extended the result to cylindrical boundary conditions. Temperley and Fisher [3], and Fisher [4] have also solved the free boundary case independently, although the generalization of the field-theoretic approach used in [4] to general graphs is less apparent.

While the solutions under various boundary conditions yield the same per-site generating function in the bulk limit, their finite-size corrections are nevertheless different. In this context there exists two unique surface structures, namely, the Möbius strip and the Klein bottle, which have escaped heretofore attention. The Möbius strip is a non-orientable surface which has a *single* side and a *single* boundary edge, and the Klein bottle, also non-orientable, has a single side and *no* edges. In view of the increasing interest on finite-size corrections, particularly their interplay with the conformal field theory [5], it is of pertinent interest to study lattice models on these surfaces. In this *Letter* we consider this problem. We solve exactly the dimer statistics for a simple quartic net embedded on a Möbius strip and on a Klein bottle. We also analyze its finite-size corrections and establish two identities relating generating functions under the Möbius and cylindrical boundary conditions.

2 Formulation

Consider first a $2M \times 2N$ simple quartic lattice $\mathcal{L}$ embedded on a Möbius strip. Here, for convenience, we assume the number of sites be even in each direction. The Möbius strip of length $2N$ and width $2M$ is shown in Fig. 1, where repeated sites are denoted by open circles. We consider close-packed dimers on $\mathcal{L}$.

Let the dimer weights be z_1 and z_2 , respectively, for dimers in the horizontal and vertical directions. We need to evaluate the dimer generating




Figure 1: A Möbius strip of a simple quartic net of length $2N$ and width $2M$. Sites denoted by open circles are repeated as indicated by the labeling. The edge orientation satisfies the Kasteleyn clockwise-odd rule for superposition polygons, and rectangles of dotted lines denote dimer cities.

function

$$Z_{(2M,2N)}^{Mob}(z_1, z_2) = \sum_{\text{dimer config.}} z_1^{n_h} z_2^{n_v}, \quad (1)$$

where n_h and n_v are, respectively, the numbers of horizontal and vertical dimers subject to the sum rule $n_h + n_v = 2MN$. The summation in (1) is taken over all close-packed dimer configurations.

A key element in the Pfaffian formulation of the generating function is to orient edges of $\mathcal{L}$ such that every superposition polygon, formed by superimposing two dimer configurations, must contain an odd number of arrows in the clockwise direction, the *clockwise-odd* rule. Kasteleyn [1] has shown that when this rule is satisfied, then the generating function of close-packed dimers can be written as a Pfaffian. For the Möbius strip $\mathcal{L}$, it can be shown that the clockwise-odd rule is satisfied if we adopt the edge orientations shown in Fig. 1. Then, we have the result

$$Z_{(2M,2N)}^{Mob}(z_1, z_2) = \text{Pf}(A) = \sqrt{\det|A|}, \quad (2)$$

where A is a $4MN \times 4MN$ antisymmetric matrix whose elements can be read off from Fig. 1.

To see the general structure of A , it is instructive to first write down the matrix A in a simpler case. For this purpose consider the 2×4 Möbius strip

($M = 1, N = 2$) shown in Fig. 2. Reading off from Fig. 2, we obtain

$$A = \begin{pmatrix} 0 & z_1 & 0 & 0 & -z_2 & 0 & 0 & -z_1 \\ -z_1 & 0 & z_1 & 0 & 0 & z_2 & 0 & 0 \\ 0 & -z_1 & 0 & z_1 & 0 & 0 & -z_2 & 0 \\ 0 & 0 & -z_1 & 0 & -z_1 & 0 & 0 & z_2 \\ z_2 & 0 & 0 & z_1 & 0 & z_1 & 0 & 0 \\ 0 & -z_2 & 0 & 0 & -z_1 & 0 & z_1 & 0 \\ 0 & 0 & z_2 & 0 & 0 & -z_1 & 0 & z_1 \\ z_1 & 0 & 0 & -z_2 & 0 & 0 & -z_1 & 0 \end{pmatrix} \quad (3)$$

and hence

$$Z_{(2,4)}^{Mob}(z_1, z_2) = 2 z_1^4 + 4 z_1^2 z_2^2 + z_2^4, \quad (4)$$

a result which can be verified by explicit enumerations.




Figure 2: A 2×4 Möbius strip.

To write down A for general M and N , we adopt “dimer cities” formed by pairing sites $\{2n, 2n + 1\}, n = 1, 2, \dots, 2MN$ as indicated in Fig. 1, and introduce 2×2 matrices

$$\begin{aligned} a(0, 0) &= \begin{pmatrix} 0 & z_1 \\ -z_1 & 0 \end{pmatrix}, & a(0, 1) &= \begin{pmatrix} -z_2 & 0 \\ 0 & z_2 \end{pmatrix} \\ a(1, 0) &= \begin{pmatrix} 0 & 0 \\ z_1 & 0 \end{pmatrix}, & a(-1, 0) &= \begin{pmatrix} 0 & -z_1 \\ 0 & 0 \end{pmatrix}. \end{aligned} \quad (5)$$

Further introduce $N \times N$ matrices

$$F_N = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, K_N = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad (6)$$

and the $2M \times 2M$ matrix

$$J_{2M} = \begin{pmatrix} & & & & 1 \\ & 0 & & & -1 \\ & & & 1 & \\ & & -1 & & \\ \dots & & & & \\ 1 & & & & 0 \\ -1 & & & & \end{pmatrix}. \quad (7)$$

Then, the matrix A can be written in terms of a linear combination of direct products of smaller ones,

$$A = A_{2N} \otimes I_{2M} + a(0, 1) \otimes I_N \otimes [F_{2M} - F_{2M}^T] + B_{2N} \otimes J_{2M}, \quad (8)$$

where

$$\begin{aligned} A_{2N} &= a(0, 0) \otimes I_N + a(1, 0) \otimes F_N + a(-1, 0) \otimes F_N^T \\ B_{2N} &= -a(1, 0) \otimes K_N + a(-1, 0) \otimes K_N^T, \end{aligned} \quad (9)$$

F_N^T and K_N^T are transposes of F_N and K_N , and I_N is the $N \times N$ identity matrix. Note that the first two terms in (8) are the corresponding A matrix for free boundary conditions.

We next consider a Klein bottle constructed by connecting the upper and lower edges of the Möbius strip of Fig. 1 in a periodic fashion with $2N$ extra vertical edges. The clockwise-odd rule again holds if these edges are oriented inversely to those in the same column. Thus, we have the matrix

$$A^{Klein} = A - a(0, 1) \otimes I_N \otimes (K_{2M} - K_{2M}^T). \quad (10)$$

3 Evaluation of determinants

We first evaluate $\det|A|$. Since J_{2M} commutes with $F_{2M} - F_{2M}^T$, and as a result they can be diagonalized by applying a common similarity transformation. Introducing the $2M \times 2M$ matrix U whose elements are

$$\begin{aligned} U_{m,m'} &= i^m \sqrt{\frac{2}{2M+1}} \sin\left(\frac{mm'\pi}{2M+1}\right) \\ (U^{-1})_{m,m'} &= (-i)^{m'} \sqrt{\frac{2}{2M+1}} \sin\left(\frac{mm'\pi}{2M+1}\right), \\ &\quad m, m' = 1, 2, \dots, 2M, \end{aligned} \quad (11)$$

we find

$$\begin{aligned} (U^{-1}I_{2M}U)_{m,m'} &= \delta_{m,m'} \\ (U^{-1}J_{2M}U)_{m,m'} &= i(-1)^{M+m} \delta_{m,m'} \\ (U(F_{2M} - F_{2M}^T)U^{-1})_{m,m'} &= (2i \cos \phi_m) \delta_{m,m'}, \quad m = 1, 2, \dots, 2M, \end{aligned} \quad (12)$$

where

$$\phi_m = \frac{m\pi}{(2M+1)}. \quad (13)$$

Thus, we can replace the $2M \times 2M$ matrices in (8) by their respective eigenvalues, and express $\det|A|$ as a product of the replaced determinants, namely,

$$\det|A| = \prod_{m=1}^{2M} \det \left| A_{2N} + (2i \cos \phi_m) a(0, 1) \otimes I_N + (-1)^{M+m} i B_{2N} \right|. \quad (14)$$

Introducing (9) and the $N \times N$ matrix

$$T_N = F_N + i(-1)^{M+m+1} K_N = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ i(-1)^{M+m+1} & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad (15)$$

we can rewrite (14) as

$$\det|A| = \prod_{m=1}^{2M} \det \left| [a(0, 0) + (2i \cos \phi_m) a(0, 1)] \otimes I_N + a(1, 0) \otimes T_N + a(-1, 0) \otimes T_N^\dagger \right|, \quad (16)$$

where $T_N^\dagger$ is the Hermitian conjugate of T_N .

Now T_N and $T_N^\dagger$ commute and they can be simultaneously diagonalized with respective eigenvalues $e^{i\theta_n}$ and $e^{-i\theta_n}$, where

$$\theta_n = (-1)^{M+m+1} \left[\frac{(4n-1)\pi}{2N} \right], \quad n = 1, 2, \dots, N. \quad (17)$$

It follows that we have

$$\begin{aligned} Z_{(2M, 2N)}^{Mob}(z_1, z_2) &= \sqrt{\det|A|} \\ &= \prod_{m=1}^{2M} \prod_{n=1}^N \left[\det \left| a(0, 0) + (2i \cos \phi_m) a(0, 1) \right| \right] \end{aligned}$$

$$\begin{aligned}
& + a(1, 0)e^{i\theta_n} + a(-1, 0)e^{-i\theta_n} \Big]^{1/2} \\
& = \prod_{m=1}^M \prod_{n=1}^N \left[4z_1^2 \sin^2 \frac{(4n-1)\pi}{4N} + 4z_2^2 \cos^2 \frac{m\pi}{2M+1} \right], \quad (18)
\end{aligned}$$

where we have used $\cos^2 \phi_m = \cos^2 \phi_{2M-m+1}$ to effect the square root. It is readily verified that (18) reduces to (4) when $M = 1, N = 2$.

The determinant $\det[A^{Klein}]$ of the matrix (10) can be evaluated in a similar fashion from which we obtain

$$Z_{(2M, 2N)}^{Klein}(z_1, z_2) = \prod_{m=1}^M \prod_{n=1}^N \left[4z_1^2 \sin^2 \frac{(4n-1)\pi}{4N} + 4z_2^2 \sin^2 \frac{(2m-1)\pi}{2M} \right]. \quad (19)$$

As a comparison, the generating functions for $2M \times 2N$ lattices with free boundaries [1] and cylindrical boundary conditions (periodic in the N direction) [2] are, respectively,

$$\begin{aligned}
Z_{(2M, 2N)}^{free}(z_1, z_2) &= \prod_{m=1}^M \prod_{n=1}^N \left[4z_1^2 \cos^2 \frac{n\pi}{2N+1} + 4z_2^2 \cos^2 \frac{m\pi}{2M+1} \right], \\
Z_{(2M, 2N)}^{cyl}(z_1, z_2) &= \prod_{m=1}^M \prod_{n=1}^N \left[4z_1^2 \sin^2 \frac{(2n-1)\pi}{2N} + 4z_2^2 \cos^2 \frac{m\pi}{2M+1} \right] \quad (20)
\end{aligned}$$

The similarity between expressions (18) - (20) is striking, with the only difference being the trigonometric factors which can be associated to the respective boundary conditions. We remark that for $(2M-1) \times 2N$ lattices we have [2]

$$Z_{(2M-1, 2N)}^{cyl}(z_1, z_2) = \frac{1}{2z_1^N} \prod_{m=1}^M \prod_{n=1}^N \left[4z_1^2 \sin^2 \frac{(2n-1)\pi}{2N} + 4z_2^2 \cos^2 \frac{m\pi}{2M} \right]. \quad (21)$$

These expressions give rise to the same bulk “free energy”

$$\begin{aligned}
f_{\text{bulk}} &= \lim_{M, N \rightarrow \infty} (4MN)^{-1} \ln Z_{(2M, 2N)}(z_1, z_2) \\
&= \frac{1}{16\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\phi \ln(4z_1^2 \sin^2 \theta + 4z_2^2 \cos^2 \phi). \quad (22)
\end{aligned}$$

Particularly, the number of ways $W_{8 \times 8}$ that an 8×8 checkerboard can be covered by 32 dominoes is obtained by setting $M = N = 4$ and $z_1 = z_2 = 1$,

yielding

$$\begin{aligned}
W_{(8,8)}^{free} &= 12\,988\,816 = 2^4 \times (901)^2 \\
W_{(8,8)}^{Mob} &= 46\,029\,729 = 47 \times 271 \times 3617 \\
W_{(8,8)}^{cyl} &= 71\,385\,601 = (8449)^2 \\
W_{(8,8)}^{Klein} &= 220\,581\,904 = 2^4 \times (3713)^2 \\
W_{(8,8)}^{toroidal} &= 311\,853\,312 = 2^8 \times 3^2 \times 135353,
\end{aligned} \tag{23}$$

where the number for toroidal boundary conditions is computed using the result of [1].

Two identities: There exist curious identities connecting generating functions for Möbius strips and cylinders. Brankov and Priezzhev [10] have proposed the identity

$$Z_{(2M,4N)}^{cyl}(z_1, z_2) = \left[Z_{(2M,2N)}^{Mob}(z_1, z_2) \right]^2, \tag{24}$$

which they established up to the order of c_3 in the large M, N expansion (26). The identity (24) can now be rigorously established using (18) and (20). The proof follows from the fact that in the product over $n = 1, 2, \dots, 4N$ on the LHS, the factor $\sin^2[(2n-1)\pi/4N]$ reproduces the set $\sin^2[(4n'-1)\pi/4N]$, $n' = 1, 2, \dots, N$, exactly twice. For $(2M-1) \times 2N$ lattices we have instead the identity (see *Note added* below)

$$Z_{(2M-1,4N)}^{cyl}(z_1, z_2) = \frac{1}{2} \left[Z_{(2M-1,2N)}^{Mob}(z_1, z_2) \right]^2. \tag{25}$$

4 Finite-size corrections

For *large* M and N , we expect to have

$$\ln Z_{(2M,2N)} = 4MN f_{\text{bulk}} + 2Nc_1 + 2Mc_2 + c_3 + O(1/N), \tag{26}$$

where c_1 , c_2 and c_3 are constants dependent at most on

$$\tau = z_2/z_1 \quad \text{and} \quad \xi = N/M, \tag{27}$$

the aspect ratio. The computation of finite-size corrections for products of the form of (18) is standard [4, 6, 7], and is outlined in the following.

We first take out a factor z_1^{2MN} in (18) and carry out the product over n by introducing the identity [8]

$$\prod_{n=1}^N \left[x^2 - 2 \cos \left(\frac{2n\pi}{N} - \alpha \right) + x^{-2} \right] = x^{2N} - 2 \cos N\alpha + x^{-2N}, \quad (28)$$

with $\alpha = \pi/2N$. This leads to

$$Z_{(2M,2N)}^{Mob}(z_1, z_2) = z_1^{2MN} \prod_{m=1}^M \left[x^{2N}(\tau, \phi_m) + x^{-2N}(\tau, \phi_m) \right], \quad (29)$$

where

$$x^2(\tau, \phi) + x^{-2}(\tau, \phi) = 2 + 4\tau^2 \cos^2 \phi, \quad (30)$$

or, explicitly,

$$x(\tau, \phi) = \tau \cos \phi + \sqrt{1 + \tau^2 \cos^2 \phi}. \quad (31)$$

It follows that we have

$$\ln Z_{(2M,2N)}^{Mob} = 2MN \ln z_1 + 2N \sum_{m=1}^M \ln x(\tau, \phi_m) + \sum_{m=1}^M \ln \left[1 + x^{-4N}(\tau, \phi_m) \right]. \quad (32)$$

We next carry out the first summation in (32) using the Euler-MacLaurin summation formula

$$\begin{aligned} \sum_{m=1}^M f(a + m\delta) &= \frac{1}{\delta} \int_a^{a+M\delta} f(\phi) d\phi + \frac{1}{2} \left[f(a + M\delta) - f(a) \right] \\ &\quad + \frac{\delta}{12} \left[f'(a + M\delta) - f'(a) \right] + O(\delta^3), \end{aligned} \quad (33)$$

with $a = 0$ and $\delta = \pi/(2M+1)$. The first term and the leading contribution of the second term lead to the following expressions for the bulk free energy,

$$\begin{aligned} f_{\text{bulk}} &= \frac{1}{2} \ln z_1 + \frac{1}{\pi} \int_0^{\pi/2} \ln x(\tau, \phi) d\phi \\ &= \frac{1}{2} \ln z_2 + \frac{1}{\pi} \int_0^{\pi/2} \ln x(\tau^{-1}, \phi) d\phi, \end{aligned} \quad (34)$$

where the second line is written down by symmetry.

After some manipulation following [6], this leads to the following for the Möbius boundary condition:

$$\begin{aligned}
c_1^{Mob}(\tau) &= \frac{1}{\pi} \int_0^{\pi/2} \ln \left[\frac{x(\tau, \phi)}{x(\tau, 0)} \right] d\phi < 0 \\
c_2^{Mob}(\tau) &= 0 \\
c_3^{Mob}(\tau, \xi) &= \ln \vartheta_3(\pi/4, q) + \frac{1}{24} \pi \tau \xi - \sum_{\ell=1}^{\infty} \ln(1 - q^{2\ell}), \quad (35)
\end{aligned}$$

where ϑ_3 is the theta function

$$\vartheta_3(u, q) = 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \cos(2nu), \quad (36)$$

with $q = e^{-\pi\xi\tau/2}$. Note that $c_2^{Mob} = 0$ is a consequence of the fact that there is no boundary edge in the M direction.

For the cylindrical boundary condition periodic in the N direction, one has in place of (29),

$$Z_{(2M, 2N)}^{cyl}(z_1, z_2) = z_1^{2MN} \prod_{m=1}^M \left[x^{2N}(\tau, \phi_m) + 2 + x^{-2N}(\tau, \phi_m) \right], \quad (37)$$

from which one obtains

$$\begin{aligned}
c_1^{cyl}(\tau) &= c_1^{Mob}(\tau), & c_2^{cyl}(\tau) &= 0 \\
c_3^{cyl}(\tau, \xi) &= c_3^{Mob}(\tau, \xi) + \ln[\vartheta_3(0, q)/\vartheta_3(\pi/4, q)]. \quad (38)
\end{aligned}$$

It follows that, for large M, N , we have

$$\frac{Z_{(2M, 2N)}^{Mob}(z_1, z_2)}{Z_{(2M, 2N)}^{cyl}(z_1, z_2)} = \frac{\vartheta_3(\frac{\pi}{4}, q)}{\vartheta_3(0, q)}. \quad (39)$$

Particularly, for $M = N$ and $z_1 = z_2 = 1$, we have $q = e^{-\pi/2}$ and (39) gives the ratio of the numbers of dimer configurations under the two respective boundary conditions as 0.701845.

For free boundary conditions we obtain

$$\begin{aligned}
c_1^{free}(\tau) &= c_1^{Mob}(\tau), & c_2^{free}(\tau) &= c_1^{Mob}(\tau^{-1}) \\
c_3^{free}(\tau, \xi) &= c_3^{Mob}(\tau, \xi) + c_1^{Mob}(\tau) + \frac{1}{2} \ln \left[\frac{2\sqrt{1+\tau^2}}{1+\sqrt{1+\tau^2}} \right]. \quad (40)
\end{aligned}$$

Here, use has been made of the second line of (34). These expressions reproduce those obtained by Fisher [4] and Ferdinand [7]. Similarly we have for the Klein bottle

$$\begin{aligned} c_1^{Klein}(\tau) &= c_2^{Klein}(\tau) = 0 \\ c_3^{Klein}(\tau, \xi) &= c_3^{Mob}(\tau, \xi) + \pi\tau\xi/8. \end{aligned} \quad (41)$$

If one takes the limit of $N \rightarrow \infty$ ($M \rightarrow \infty$) first while keeping M (N) finite [5], one obtains from (26)

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{2N} Z_{(2M, 2N)} &= 2M f_{bulk} + c_1 + \Delta_1/2M + O(1/M^2), \\ \lim_{M \rightarrow \infty} \frac{1}{2M} Z_{(2M, 2N)} &= 2N f_{bulk} + c_2 + \Delta_2/2N + O(1/N^2). \end{aligned} \quad (42)$$

One finds after some algebra the results

	Möbius	Klein	Free	Cylindrical	Toroidal
c_1	$c_1^{Mob}(\tau)$	0	$c_1^{Mob}(\tau)$	$c_1^{Mob}(\tau)$	0
c_2	0	0	$c_1^{Mob}(1/\tau)$	0	0
Δ_1	$\pi\tau/24$	$\pi\tau/6$	$\pi\tau/24$	$\pi\tau/24$	$\pi\tau/6$
Δ_2	$\pi/24\tau$	$\pi/24\tau$	$\pi/24\tau$	$\pi/6\tau$	$\pi/6\tau$

where we have also included results for toroidal boundary conditions worked out from results of [7]. The tabulated values of Δ_1 and Δ_2 are consistent with the finding of [5] that the periodical boundary condition (in the lateral direction of an infinite strip) is distinct from the free or fixed, now including the Möbius, boundary conditions. While it is tempting to extract the central charge c from Δ_1 and Δ_2 resulting in $c = \tau$ and τ^{-1} , respectively, we remark that the extraction is superfluous since results of [5] apply to systems at criticality while the present dimer system does not exhibit a critical point.

Note added: After the submission of this paper we learned of a recent preprint by Tesler [9] in which the dimer generating function $Z_{(\mathcal{M}, \mathcal{N})}^{Mob}(z_1, z_2)$ is enumerated for all $\mathcal{M}, \mathcal{N}$ (even or odd) as linear combinations of two Pfaffians with the results given in terms of a q -analogue of the Fibonacci number. We have verified that for $\mathcal{M}, \mathcal{N}$ even Tesler's result is the same as our expression (18). For $\mathcal{M}$ odd and $\mathcal{N}$ even, Tesler found the solution also given by (18) but with an extra factor $z_1^{-\mathcal{N}/2}$ with $2M \rightarrow \mathcal{M}$, $2N \rightarrow \mathcal{N}$, and the product over m taken from 1 to $(\mathcal{M} + 1)/2$. We remark that this

leads to the identity (25) upon using (21). For $\mathcal{M}$ even and $\mathcal{N}$ odd, however, the solution assumes a more complicated expression [9]. The Möbius strip problem has also been studied by Brankov and Priezzhev [10] in the context of the large M, N expansion (26) of the free energy.

5 Acknowledgement

We are grateful to R. P. Stanley for calling our attention to Ref. [9] and V. B. Priezzhev for pointing out to us Ref. [10]. We are also indebted to P. Kleban for an enlightening discussion on the connection with the conformal field theory. This work has been supported in part by NSF Grant DMR-9614170.

References

- [1] P. W. Kasteleyn, *Physica*, 27 (1961) 1209.
- [2] For a comprehensive review on dimer statistics using Pfaffians see B. M. McCoy and T. T. Wu, *Two-dimensional Ising model* (Harvard University Press, Cambridge, Massachusetts, 1973)
- [3] H. N. V. Temperley and M. E. Fisher, *Phil. Mag.* 6 (1961) 1061.
- [4] M. E. Fisher, *Phys. Rev.* 124 (1961) 1664.
- [5] H. W. J. Blöte, J. L. Cardy and M. P. Nightingale, *Phys. Rev. Lett.* 56 (1986) 742.
- [6] M. N. Barber and B. W. Ninham, *Random and restricted walks*, (Gordon and Breach, New York 1970), Appendix AI.
- [7] A. D. Ferdinand, *J. Math. Phys.* 8 (1967) 2332.
- [8] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, (Academic Press, New York 1994) 1.394.
- [9] G. Tesler, *Matchings in graphs on non-oriented surfaces*, preprint.
- [10] J. G. Brankov and V. B. Priezzhev, *Nucl. Phys.* B400(FS) (1993) 633.